BIRD IN THE HAND.

H.J.de S.Disney, Australian Museum, Sydney. Drawings by B.Bertram, Australian Museum Art Dept.

The Striated and Brown Thornbills.

Although these birds may be distinguished by their different calls and habits in the field, when trapped some confusion may arise over distinguishing them, since both kinds are about the same size and brown with lighter underparts striped black. No attempt is made here to distinguish the races of the Brown and most observations have been made from the form found in eastern N.S.W. The descriptions and drawings only give diagnostic characters.

STRIATED THORNBILL Acanthiza lineata R.A.O.U. No.470

Adult:

Crown.

Brown with narrow streaks formed by · white shafts of crown feathers.


Iris: Dark Grey


Sub-Adult:


Similar to adult, and no satisfactory difference yet observed.

Juvenile:

Similar to adult, but throat and breast markings indistinctly mottled rather than striped.







Adult:

Forehead with crescent shaped or "scalloped" light brown edges to feathers.

Throat & Breast. Greyish white streaked black, in race katherina (N.E.Qld.) streaks faint similar to juvenile of eastern race.

Belly. Greyish.

Iris. Full red brown.

Sub-Adult: Similar to adult except:

Iris. With chestnut ring round pupil surrounded by olive grey. As bird matures iris becomes full red brown with no outer grey.

Juvenile:

<u>Forehead</u>. With faint scalloping.

Throat & Breast. Buff grey with faint streaks.

<u>Iris</u>. Chestnut brown with grey outside ring.

Some races of pusilla have well marked chestnut upper tail coverts.

CORRIGENDUM

The scientific names given in Bird in the Hand Vol.2 No.1 p.16 were taken from an unamended check list. These names have been altered and should read as follows:-

Fantailed Cuckoo RAOU No. 338

Cacomantis pyrrhophanus

Brush (Square-tailed) Cuckoo RAOU No. 339

Cacomantis variolosus