



## I N D E X

	Page
<i>Some Retrap Data On The Brown Thornbill</i> .....	3
<i>Mass Movement of Honeyeaters</i> .....	8
<i>Grey-Crowned Babblers</i> .....	8
<i>Backyard Trapping</i> .....	9
<i>The Silver Gull Banding Programme</i> .....	14
<i>More "Senior" Silvereyes</i> .....	14
<i>A Study Of Birds In The Antarctic and Subantarctic</i> .....	15
<i>Bird In The Hand</i> .....	17
<i>Recovery Round-up</i> .....	18
<i>Queensland-banded Swamphen Recovered From West Irian</i> .....	19
<i>Annual General Meeting and Scientific Meeting</i> .....	20
<i>New Members</i> .....	21
<i>Editorial</i> .....	22
<i>Extracts From Letters</i> .....	23

### **Cover Picture —**

Silver-grey Fulmar (*Fulmarus glacialisoides*) at Ardery Island, Wilkes, Antarctica. This species may be seen in high altitudes throughout the Southern Ocean during the winter; in October the birds return to their breeding colonies before the snow has melted. They burrow down through the snow to their nesting ledges of the previous season. Recently, a flea has been discovered adapted to life in these circumstances (see page 16).

The statements and opinions contained in the papers published in this journal are the responsibility of the respective authors.

**All Rights Reserved.**

**PRICE 5/-**

[Registered at the G.P.O. for transmission by post as a periodical.]