



• Red wattle-bird (*Anthochaera carunculata*) caught for banding.—Picture, S. G. Lane.

574 Eastern Silvereye (*Zosterops lateralis*)

- (a) 010-78709. Adult banded by R. Curran at Pendle Hill (Sydney) N.S.W. on 20.5.64. Retrapped at banding place on 22.5.64 and 4.6.64 then recovered (dead) at St. Mary's Tas. about 25.11.64. 560 miles S.S.W.
 (b) 010-98280. Adult banded by J. Liddy near Cudgen, N.S.W. on 15.6.64. Recovered at

Silvan, Vic. about 15.11.64. 823 miles S.W. (This is the most distant recovery to date.)

575 Western Silvereye (*Zosterops lateralis gouldi*)

010-35847. Banded by R. H. Stranger at Nedlands, W.A. on 8.2.64. Recovered at Yokine, W.A. on 6.3.65. 6 miles N.N.E.

630 Crescent Honeyeater (*Phylidonyris pyrrhoptera*)

021-03150 Banded by D. J. Noonan on 21.1.65 at Ben Cairn (between Healesville and Warburton, Vic.) Retrapped by bander at Panton's Gap, Vic. on 19.2.65. Released with band. 3 miles W.N.W. and 1,700 ft. below banding place.

638 Red Wattle-bird (*Anthochaera carunculata*)

070-47025 Juvenile banded by J. P. Ipsen, 17 miles N. of Bendigo, Vic. on 24.3.64. Found dead at Ringwood, Vic. in January, 1965. 108 miles S.S.E.

690 Raven (*Corvus coronoides*)

100-41974. Adult banded by E. B. Thomas, 153 miles W. of Barham, N.S.W. on 29.8.64. Recovered at Thornton, Vic. on 9.3.65. 160 miles S.E.

999 Common Starling (*Sturnus vulgaris*)

- (a) 060-16429 Banded by R. H. Green at Launceston, Tas., on 4.5.64. Recovered at Oatlands, Tas., on 22.11.64. 60 miles S.
 (b) 060-16781. Banded by R. H. Green at Launceston, Tas., on 5.5.64. Recovered at Jericho, Tas., between 7th and 19th October, 1964. 66 miles S.

RECOVERY DETAILS

Recovery details are obtained mainly from two sources—

1. The Australian Bird-banding Scheme;
2. Individual banders.

The Secretary of the A.B.B.S. has kindly arranged for copies of interesting recoveries to be forwarded to the Association. It is neither possible nor expedient for all recoveries to be made available through this source; consequently some recoveries of considerable value may not be included in Recovery Round-up unless individual banders provide the details.

Recoveries 10 miles from banding place for some species considered sedentary may be of greater value than, for instance, a more spectacular shearwater recovery of 5,000 miles.

Banders are requested to forward to me (or the Hon. Editor) details of any recovery or re-trap that they consider may be suitable for publication. The editor will finally determine which ones will be included.

In order to avoid possible errors, the "Report to Bander" form should be forwarded. If the bander requires the form to be returned later, the top of the form should be clearly endorsed to this effect. The enclosure of a stamped, addressed envelope would be appreciated.

—S. G. Lane, 65 Wood St., Lane Cove, N.S.W.

Subscriptions

Subscriptions are now overdue. **No further issues** will be forwarded to unfinancial members. In order to maintain and improve the standard of the journal, the Committee requests the co-operation and support of all members in an endeavour to increase membership. Any donations will be welcome.