

- Klages, N. T. W., Pemberton, D. and Gales, R. P. (1990). The diets of King and Gentoo Penguins at Heard Island. *Aust. Wildl. Res.* 17: 53–60.
- LaCock, G. D., Hecht, T. and Klages, N. (1984). The winter diet of Gentoo Penguins at Marion Island. *Ostrich* 55: 188–191.
- Mauchline, J. (1980). Measurement of body length of *Euphausia superba* Dana. *Biomass Handbook* 4: 1–9.
- Patterson, G. B. (1986). A statistical method of testing for dietary differences. *N.Z. J. Ecol.* 13: 113–115.
- Ridoux, V. (1988). Subantarctic krill, *Euphausia vallentini* Stebbing, preyed upon by penguins around Crozet Island (Southern Indian Ocean): population structure and annual cycle. *J. Plank. Res.* 10: 675–690.
- Volkman, N. J., Presler, P. and Trivelpiece, W. (1980). Diets of pygoscelid penguins at King George Island, Antarctica. *Condor* 82: 373–378.
- Williams, A. J. (1980). Aspects of the breeding biology of the Gentoo Penguin, *Pygoscelis papua*. *Gerfaut* 70: 283–295.
- Williams, R. and McEldowney, A. (1990). A guide to the fish otoliths from waters off the Australian Antarctic Territory, Heard and Macquarie Islands. ANARE Research Notes 75.
- Williams, T. D. (1991). Foraging ecology and diet of Gentoo Penguins *Pygoscelis papua* at South Georgia during winter and an assessment of their winter prey consumption. *Ibis* 133: 3–13.

Corella, 1992, 16(5): 132–133

LONGEVITY OF CAPE PETRELS *Daption capense* AT BLUFF ISLAND, VESTFOLD HILLS, EAST ANTARCTICA

F. I. NORMAN and S. J. WARD¹

Department of Ecology and Evolutionary Biology, Monash University, Clayton, Victoria 3168.
¹Present address: Department of Zoology, James Cook University, Townsville, Queensland 4811.

Received 24 September, 1991

The Cape Petrel *Daption capense* nests on the Antarctic Peninsula, on Antarctic and Subantarctic islands as well as on the continent itself (Watson 1975; Marchant and Higgins 1990). Breeding occurs in East Antarctica, including sites in Prydz Bay (Woehler and Johnstone 1991). Bluff Island (68°33'S, 77°54'E), off Davis Station in the Vestfold Hills, is one such nesting area where Cape Petrels have been banded intermittently since 1959. To the end of the 1989/90 summer 299 pulli and 130 adult birds at nests have been marked there.

Bluff Island was visited briefly in early February 1989 and again, for a longer period, on 7 January 1991 during studies on the South Polar Skua *Catharacta maccormicki*. Previously-banded Cape Petrels were noted in 1989. In 1991 petrels at nests over much of the Island were examined, particularly where nest density was high and where birds banded prior to 1989 were breeding. Of 334 adults caught at nest sites, 15 had been banded previously. Four adults were banded in 1989, one breeding adult was banded as a chick in February 1974 and another as a pullus in 1984;

a further adult had been banded as a nestling at the Forbes Glacier (67°38'S, 62°21'E), some 800 km away near Mawson in February 1973. Eight adults were banded on nests in February 1984, when 68 pulli and 32 breeding birds were marked by K. Green.

Since Cape Petrels are essentially monogamous, with a long-lasting pair bond, and show a strong fidelity to nesting areas (Marchant and Higgins 1990), such recapture data may provide a measure of minimum survival and hence annual mortality. Minimum mean annual survival (S) may be calculated as $S = (r/n)^{1/y}$ or, as a percentage, as $S = 100 \cdot \sqrt[y]{r/n}$, where y = number of years, n = number originally banded, and r = number of recaptures (e.g. Mougín 1975). Thus for the cohort of adults marked in 1984, minimum mean survival was 0.82 (82%) and hence annual mortality (m) 0.18 (18%). Life expectancy (E) = $2 - m/2m$ ($200 - m/2m$ in percentage form) and, in consequence, is some five years for such adult birds assuming that mortality is independent of age. Confidence limits about such estimates are, however, large; mortality estimates from these data, for example, vary between 12 and 28 per cent.

Conclusions from these data are limited both by the small numbers of banded birds in the original cohort and their subsequent recaptures. In addition, mortality estimates may be magnified by band loss (presumably minimized by the use of stainless steel bands from 1984 onwards), the non-recapture of previously-marked birds (reduced here by the coverage of the Island in areas of previous bandings) and emigration. However, Mougín (1975) reported some 21 per cent of marked birds at the same nest sites six years later and in this study at least 25 per cent of adult birds banded on Bluff Island in 1984 were present there in 1991, seven years later. Estimated mortality rates are, however, somewhat higher than 4–7 per cent previously reported (Hudson 1966; Mougín 1975), and life expectancy correspondingly less. That only one of 68 chicks banded in 1984 was recaptured as a breeding adult could suggest heavy mortality of such chicks that season (or subsequent prebreeding mortality). However, since the majority of birds breed for the first time at about five years of age (Pinder 1966; Beck 1969) and hence would have been available for capture in 1991, it may also indicate some dispersal to breed at other sites (e.g. Beck 1969), rather

than strict philopatry (e.g. Pinder 1966; Marchant and Higgins 1990). The recapture of an adult Cape Petrel at Bluff Island originally banded (with a monel band) as a chick near Mawson indicates that some birds do disperse from natal sites to breed elsewhere, sometimes over considerable distance. Pinder (1966) noted that 5–16 per cent of the breeding population was replaced in any one year and hence estimated adult mortality for Cape Petrels at Bluff Island may reflect both death and emigration. Such estimates may be biased further by failure of nesting attempts early in the breeding season, and the subsequent dispersal of adults. At breeding colonies such as Bluff Island, blizzards and resulting snow drifts could reduce nest site availability, nesting attempts and success, and lead to abandonment of nesting colonies. Further studies may clarify such issues.

ACKNOWLEDGMENTS

We thank the Director, Australian Antarctic Division for the opportunity to visit Bluff Island, and to the Station Leader and fellow expeditioners at Davis during the 1990/1991 summer for their assistance in many ways. Annie Wessing helped in the examination of the Bluff Island birds and J. M. Cullen, W. B. Emison and anonymous referees commented helpfully on the manuscript. We are grateful too to the previous ANARE expeditioners who banded Cape Petrels on Bluff Island, and thank the Australian Bird and Bat Banding Scheme who provided the appropriate details.

REFERENCES

- Beck, J. R. (1969). Food, moult and age of first breeding in the Cape Pigeon *Daption capensis* Linnaeus. *Brit. Antarct. Surv. Bull.* 21: 33–44.
- Hudson, R. (1966). Adult survival estimates for two antarctic petrels. *Brit. Antarct. Surv. Bull.* 8: 63–73.
- Marchant, S. and Higgins, P. (1990). (Eds). 'The Handbook of Australian, New Zealand and Antarctic birds.' Vol 1. (Oxford University Press: Melbourne.)
- Mougín, J. L. (1975). Ecologie comparée des Procellariidae antarctiques et subantarctiques. *C.N.F.R.A.* 36: 1–195.
- Pinder, R. (1966). The Cape Pigeon, *Daption capensis* Linnaeus, at Signy Island, South Orkney Islands. *Brit. Antarct. Surv. Bull.* 8: 19–47.
- Watson, G. E. (1975). 'Birds of the Antarctic and Subantarctic.' (Amer. Geophys. Union: Washington.)
- Woehler, E. J. and Johnstone, G. W. (1991). The status and conservation of the seabirds of the Australian Antarctic Territory. In 'Seabird status and conservation: a supplement.' (Ed. J. Croxall). *ICBP Technical Publication* 11: 279–308.