
# Restless Flycatcher Mylagra inquieta Species No.: 728 Band Size: 02



### Morphometrics:

Occurs in southern and eastern Australia to the eastern half of Cape York Peninsular

	Adult Male	Adult Female
Wing:	98 – 110 mm	94 -110 mm
Tail:	86 -104 mm	87 – 100 mm
Bill (tip to skull):	21.6 – 24.5 mm	21.6 – 24.6 mm
Bill Width:	5.5 – 6.7 mm	5.8 – 6.6 mm
Tarsus:	17.5 – 20.1 mm	17.5 – 20.0 mm
Weight:	18 – 25 g	13 – 23 g

#### Ageing:

Juvenile and immature, see drawings;

First Immature – similar to adults, but retain brown alula and other wing feathers;

Adult plumage attained late first or early second year;

Breeding commences in August, so full adult plumaged birds captured post winter are aged (2+).

#### Sexing:

Adult Male Adult Female

Upperparts: mostly black, but back and rump slightly duller than male with less

grey-black all with bluish sheen; sheen;

Lores, eye-ring & black; dark grey;

ear coverts:

Breast: usually white, but may have dull usually has orange-buff flush;

orange-buff flush reducing with age;

## Similar species:

The Paperbark Flycatcher *Myiagra nana* (formerly known as the Little Restless Flycatcher) occurs in northern Australia from the Kimberley region of WA to the south-western portion of Cape York Peninsular. It is very similar in plumage to the Restless Flycatcher, but is considerably smaller – see separate profile sheet.