

- Briggs, S. V. (1977). Variation in waterbird numbers at four swamps on the northern tablelands of New South Wales. *Aust. Wildl. Res.* 4: 301-309.
- Crawford, D. N. (1979). Waterbirds: indices and fluctuations in dry season refuge areas, Northern Territory. *Aust. Wildl. Res.* 6: 97-103.
- Curry, P. (1981). A survey of the birds of Herdsman Lake, 1980-1981. Department of Conservation and Environment, Western Australia, Bulletin 105.
- Department of Science, Bureau of Meteorology (1978-1981). Monthly weather review, Western Australia.
- Ford, J. R. (1958). Seasonal variation in populations of Anatidae at the Bibra Lake District, Western Australia. *Emu* 58: 31-41.
- Frith, H. J. (1982). Waterfowl in Australia, 2nd ed. Angus and Robertson, Sydney.
- Gentilli, J., and Bekke, H. (1983). Modelling a climatically pulsating population: Grey Teal in south-western Australia. *J. Biogeogr.* 10: 75-96.
- Lamm, D. W. (1964). Seasonal counts of birds at Lake George, New South Wales. *Emu* 64: 115-128.
- Lane, J. A. K., and Munro, D. R. (1981). 1980 review of rainfall and wetlands in the south-west of Western Australia. Department of Fisheries and Wildlife, Western Australia. Report 47.
- Morris, K. D., and Knott, B. (1979). Waterfowl utilization of Lake Claremont during 1977. *West Aust. Nat.* 14: 145-149.
- Pen, L. J. (1983). Peripheral vegetation of the Swan and Canning estuaries 1981. Department of Conservation and Environment, Western Australia, and Swan River Management Authority, Bulletin 113.
- Riggert, T. L. (1977). The biology of the Mountain Duck on Rottnest Island, Western Australia. *Wildl. Monogr.* 52: 1-67.
- Sedgewick, E. H. (1973). Birds of the Benger Swamp. *West Aust. Nat.* 12: 147-155.
- Serventy, D. L. (1948). The birds of the Swan River District, Western Australia. *Emu* 47: 241-286.
- Serventy, D. L., and Whittell, H. M. (1976). Birds of Western Australia. 5th ed. University of Western Australia Press, Perth.
- Whyte, R. J. (1981). Winter fluctuations in waterbird numbers on a northern tablelands lagoon of New South Wales. *Emu* 81: 243-246.

DATA EXCHANGE

This section has been included in the Journal at the request of a number of members of the Association. It is designed to allow the inclusion of material of a brief and possibly preliminary nature with a very short publication time. Suitable subject matter would include Weights and Measurements, Moults, Plumage changes, Colour of soft parts, Ageing and Sexing methods and Requests for information, etc.

It is recommended that material from 'Data Exchange' be referenced as, e.g.:

Lane, S. G. (1983), 'Weights and Measurements — Eastern Spinebill', In Data Exchange, *Corella* 7: 22.
Hon. Editor

Weights and Measurements

Long-billed Corellas *Cacatua tenuirostris*,
collected in western Victoria 1977-1983.
Monthly Weights (g)

MALES				
Month	Range	Mean	SD	n
Jan.	519-587	554	24.5	8
Feb.	531-629	577	32.4	16
Mar.	489-676	589	40.1	3
Apr.	557-640	599	34.1	4
May	528-690	602	51.5	16
June	547-639	593	30.4	10
July	491-718	627	71.8	9
Aug.	485-611	566	45.9	7
Sep.	558-689	608	46.0	10
Oct.	536-635	586	28.2	9
Nov.	503-678	577	47.0	23
Dec.	524-659	597	36.9	20
FEMALES				
Month	Range	Mean	SD	n
Jan.	497-568	532	21.7	17
Feb.	473-617	544	41.8	21
Mar.	458-636	528	44.9	27
Apr.	467-624	538	42.1	14
May	493-608	547	38.1	7
June	535-571	559	12.4	8
July	477-620	557	49.7	11
Aug.	462-630	551	47.3	13
Sep.	505-718	551	57.9	13
Oct.	500-580	550	35.1	4
Nov.	515-615	562	32.6	24
Dec.	447-599	542	42.9	10

Most specimens were shot under permit by grain growers and then frozen until they could be picked up by staff of the Fisheries and Wildlife Service.

Sex was determined by dissection; age was not determined; the weights of a few birds (in November and December) which obviously had just fledged were not included.

W. B. Emison and I. D. Temby,
Arthur Rylah Institute for Environmental Research,
Fisheries and Wildlife Service,
123 Brown Street,
Heidelberg, Vic. 3084.