

population. I found, as did Lane (1972b), that the numbers of Silvereyes present tended to vary considerably from year to year. In particular large numbers were present during 1970 to 1972.

I found trapping to be much more effective than mist-netting. Birds rapidly become net shy and avoid nets, whereas they enter traps readily and repeatedly and give a more accurate indication of birds present in the area.

ACKNOWLEDGEMENTS

I am grateful to Messrs Max Waterman and Warren Hitchcock for reading a preliminary draft and to Annette Cam, Kim Lowe and Graham Cam, who assisted in the preparation of the manuscript's final draft. I wish to acknowledge Joan Paton for the use of her plumage-colour codes.

REFERENCES

- Hitchcock, W. B. (1966). Tenth Annual Report of the Australian Bird-banding Scheme. July 1963 to June 1964. *CSIRO Wildl. Res. Tech. Paper* No. 11.
- Lane, S. G. (1966). The Co-operative Silvereye Project. *Aust. Bird Bander* 4: 73-75.
- Lane, S. G. (1972a). A Review of the Co-operative Silvereye Project. *Aust. Bird Bander* 10: 3-6.
- Lane, S. G. (1972b). Tasmanian Type Silvereyes in New South Wales. *Aust. Bird Bander* 10: 33-34.
- Lane, S. G. and Battam, H. (1971). Silvereye movement in eastern Australia. *Aust. Bird Bander* 9: 80-82.
- Liddy, J. (1966). A summary of Silvereye banding. *Aust. Bird Bander* 4: 71-73.
- McKean, J. L. (1965). Weights of the Silvereye at Canberra. *Aust. Bird Bander* 3: 43-44.
- Mees, G. F. (1974). The migration of the Tasmanian race of the Silvereye. *Aust. Bird Bander* 12: 51-54.
- Robertson, J. S. (1971). South-East Queensland aspects of the Co-operative Silvereye Project. *Aust. Bird Bander* 9: 51-55.
- Robertson, J. S. (1972). Silvereye colour cline. *Aust. Bird Bander* 10: 12-13.
- Swanson, N. M. (1968). Suburban Silvereyes. *Aust. Bird Bander* 6: 5-7.
- Walker, D. M. (1962). Three years Silvereye banding—an analysis. *Aust. Bird Bander* 1: 26-29.
- Walker, D. M. (1964). The relationship between plumage colouration, breeding location and bodyweight in the Silvereye (*Zosterops lateralis*). *Aust. Bird Bander* 2: 79-81.

Corella, 1987, 11(2): 51

THE DIET OF WEDGE-TAILED EAGLES *Aquila audax*

In Cindy Hull's recent article (Hull 1986) on this subject, based on a study of pellets and other prey remains at nest sites and roosting trees, I noted a list of species not previously recorded in the Wedge-tailed Eagles' diet. These species included the Sacred Ibis *Threskiornis aethiopica*. The following observations, as recorded in my diary, may therefore be of interest.

On 6 April 1980, I was motoring near Yuckandandah, Victoria, when I saw two Wedge-tailed Eagles on the ground, one standing on a strugg-

ling Sacred Ibis while the other looked on. When disturbed, the attacking eagle backed-off and the ibis got to its feet and staggered to the shelter of some nearby rushes. The eagles then flew up into a tall adjacent eucalypt, where a number of ravens had gathered. We then left the scene.

Hull, C. (1986). The diet of the Wedge-tailed Eagle, *Aquila audax*, breeding near Melbourne. *Corella* 10: 21-24.

Arthur Gwynn,
40 Golfers Parade,
Pymble, N.S.W. 2073