

The other sightings, except for the resident King Penguins, are extra-limital records that add to the knowledge of each species' pelagic dispersion.

ACKNOWLEDGEMENTS

The late Gavin Johnstone initiated recent ornithological investigations at Heard Island with his work there in 1980, and maintained that interest up to the time of his death. The assistance of the Australian Bird and Bat Banding Schemes is gratefully acknowledged. I would like to thank Max Downes for permission to use his Southern Giant-Petrel data, and all members of the 1985, 1986-87 and 1987-88 ANARE whose observations are reported here.

REFERENCES

- Allison, I. F. and Keage, P. L. (1986). Recent changes in the glaciers of Heard Island. *Polar Rec.* 23: 255-271.
- Burton, H. R. (Compiler), (1989). Heard Island ANARE 1987-88 report (unpublished).
- Burton, H. R. and Williams, D. L. (Compilers), (1986). Heard Island ANARE 1985 report (unpublished).
- Chittleborough, R. G. and Ealey, E. H. M. (1950). Bird-ringing at Heard Island during 1949. *Emu* 50: 102-104.
- Croxall, J. P. (1981). Aspects of the population demography of Antarctic and Subantarctic seabirds. *CNFERA* 51: 479-488.
- Downes, M. C., Ealey, E. H. M., Gwynn, A. M. and Young, P. S. (1959). The birds of Heard Island. *ANARE Rep. Ser. B(1)*: 135pp.
- Downes, M. C., Gwynn, A. M. and Howard, P. F. (1954). Banding of giant petrels at Heard and Macquarie Islands. *Emu* 54: 257-262.
- Howard, P. F. (1954). Banding of the Black-browed Albatross at Heard Island and Macquarie Island. *Emu* 54: 256.
- Howard, P. F. (1956). Banding of giant petrels at Heard Island and Macquarie Island - II. *Emu* 56: 401-404.
- Lambeth, A. J. (1951). Heard Island geography and glaciology. *J. Proc. R. Soc. N.S.W.* 84: 92-98.
- Ledingham, R. (Compiler), (1987). Heard Island Expedition ANARE 1986-87 report (unpublished).
- Woehler, E. J. (1989). The status and conservation of the seabirds of Heard Island and the McDonald Islands. ICBP Tech. Publication.
- Woehler, E. J. and Johnstone, G. W. (1988). Banding studies of giant petrels, *Macronectes* spp., at Macquarie Island. Proc. Macquarie Island Symp. May 1987. *Pap. Proc. R. Soc. Tasm.* 122: 143-152.

THE ANTARCTIC FLEA

Glaciopsyllus antarcticus

Bell, P. J., Burton, H. R. and van Franeker, J. A. (1988) have reported in *Polar Biology* 8: 403-410 the findings of a study on Ardery Island. The life-cycle of the flea was completed only where birds were present as the flea is dependent upon its host for food and warmth, and was completed during the incubation of eggs and rearing of chicks of the Antarctic Fulmar, before the chicks were fledged. Flea eggs were found in nest material while the fulmars were incubating eggs, and flea larvae when chicks were hatching and during their subsequent development. However, larvae were found mainly in the belly down of the chicks, where they fed on blood and/or blood-faeces from adult fleas, and here they pupated, in woven cocoons attached to the down feathers. Adults emerged before the chicks had fledged; a few were found in the nests but they were abundant on chicks (505 in the belly down of one chick). Mating adults were seen on chicks, and it appeared that fleas on chicks overfed, that is, they produced large quantities of faeces containing blood of the host. These faeces are often a source of food for larval fleas. The authors feel that this synchronization of the breeding cycle of the flea with that of its host and its dependence upon chicks, suggest it may be like the rabbit flea *Spilopsyllus cuniculi*, whose reproductive biology is controlled by the hormonal state of the host. If true, it will be the first avian example. This flea is clearly not a nest-flea like most bird fleas. The question remains — how does the flea overwinter? The lack of fleas in the nests of Antarctic Fulmars known to have been well infested before the bird's departure has led the authors to think that adult fleas remain on non-breeding hosts while they are away from their nesting sites. Clearly all Antarctic Fulmars and Snow Petrels captured alive between April and November, when they are not breeding, should be examined closely for fleas, and there is a strong case for examining carefully washed-up birds if in good condition. It only requires one infested bird to be found to indicate that the authors may be correct. Any fleas found should be preserved, preferably in 70% alcohol (a nip of whisky would be warranted), so that the flea may be identified correctly.

Durno Murray