
Magpie-lark Grallina cyanoleuca





Morphometrics:

Two subspecies – nominate G.c. cyanoleuca south of c. 20° S and G.c. neglecta north thereof. Measurements presented here are for G.c. cyanoleuca:

	Adult Male	Adult Female
Wing:	168 – 195 mm	162 – 184.5 mm
Tail:	111 – 141 mm	108 – 136 mm
Weight:	69.3 – 118 g	70 – 94.5 g

Ageing:

Both juvenile and first immatures distinguished by plumage and bare parts;

Juvenile plumage is retained only for about three months after fledging;

First immature plumage differs from adult by retaining most or all juvenile remiges, rectrices outer greater secondary coverts, greater primary coverts and feathers of alula which are contrastingly browner and more worn than in adults;

Immatures acquire adult plumage within their first year, coinciding with adult post-breeding moult which commences in Nov.-Dec and is completed by Apr.-May.;

Thus birds in complete adult plumage August to March will be aged (2+);

Adult (2+) Juvenile

Bill: cream with blackish culmen, tip to grey-black, blackish or dark grey with pale tip;

upper mandible and nasal groove;

Gape: grey-black; pale yellow; lris: pale straw-yellow; dark olive brown;

Forehead: see below; black (as for adult male); Throat: see below; white (as for adult female); Eyebrow: see below; white (as for adult male);

Sexing:

The easy way to sex adults – girls have clean faces and boys have dirty faces! Otherwise:

Adult Male Adult Female

Forehead: black; white; Throat: black; white;

Eyebrow: white black above & white behind eye;