White-throated Gerygone Gerygone olivacea Species No.: 453 Band size: 01



Variation in pattern of undertail between subspecies





G.o. olivacea (e. Australia)

Morphometrics:

Bill:

 $\begin{array}{ccccc} & \text{Adult Male} & \text{Adult Female} \\ \text{Wing:} & 55-62 \text{ mm} & 54-60 \text{ mm} \\ \text{Tail:} & 37-44 \text{ mm} & 36-43 \text{ mm} \\ \text{Weight:} & 5.5-8.5 \text{ g} & 6.0-8.4 \text{ g} \end{array}$

Ageing:

Adult (1+) Juvenile

black or dark brown; grey black upper mandible and dark

pink-brown lower mandible; with a

darker tip;

Tomia: black or dark brown; dull pinkish;

Gape: black or grey-black; dull pinkish or pale yellowish;

Iris: red, crimson or orange-red; grey-brown, light reddish brown, mid

brown or olive-brown;

Lores: dark brown; greyish-yellow or greyish-olive;

Supra-loral spot: white; yellow;

Chin & throat: yellow or pale-yellow

Juveniles are distinguished by plumage and bare parts;

Immatures resemble adults but may have patchy pale yellow feathering on the chin and throat; Immatures might also be distinguished by the presence of synchronous growth or fault bars in retained juvenile wing and tail feathers;

Immatures attain complete adult plumage when approximately one year old.

Sexing:

There is no sexual dimorphism in plumage or size;

Incubation is mostly done by the female though males incubate sometimes so both sexes will have brood patches.