


Western Yellow Robin *Eopsaltria griseogularis* Species No.: 394 Band size: 02



Morphometrics:

| | Nominate <i>E.g. griseogularis</i> | | ssp. <i>E.g. rosinae</i> | |
|---------|------------------------------------|-------------------|--------------------------|-------------------|
| | Adult Male (2+) | Adult Female (2+) | Adult Male (2+) | Adult Female (2+) |
| THL: | | | | |
| Wing: | 83 – 94 mm | 80 – 91 mm | 85 – 97 mm | 81 – 89 mm |
| Tail: | 55 – 68 mm | 57 – 66 mm | 62 – 73 mm | 58 – 66 mm |
| Weight: | 20.0 – 21.5 g | 18.5 – 20.5 g | 17.4 – 22.5 g | 17.0 – 22.0 g |

The two subspecies interbreed though an intergrade zone.

Ageing:

- All traces of mottled brown Juvenile plumage replaced with adult-like body plumage within three months of fledging;
- Juvenile grey-brown remiges and rectrices and usually some dark brown coverts with white tips are retained by Immature.
- Tail feathers of Immatures are pointed.
- Moults to adult plumage occurs early in the second year.
- Thus all adults 2+ and Immatures (1) in Autumn/Winter or (2-) in Spring.

Sexing :

Females are generally slightly smaller than males
Female alone incubates.