


White-streaked Honeyeater *Trichodere cockerelli* Species No.: 627 Band size: 03



Morphometrics:

	Adult Male	Adult Female
Wing:	74 – 81 mm	64 – 68 mm
Tail:	58 – 70 mm	53 – 59 mm
Weight:	14 – 20 g	12.0 – 13.5 g

Ageing:

	Adult (2+)	Immature (2-) or (1)	Juvenile
Bill:	nostrils to tip – black, basal half dark blue-grey, light blue-grey, pale grey or dark greenish-blue;	as for adult;	dark grey to grey-black with pale grey base;
Gape:	dark grey-blue;	dark grey-blue;	pale grey, cream or yellow
Iris:	red-brown or dark brown;	dark brown;	dark brown;
Forehead, crown nape & lores:	grey-black;	grey black;	brown;
Ear coverts:	broad yellow triangular tuft;	narrow elongated rear tuft;	indistinct small roundish yellow tuft at rear;
Chin & throat:	pale grey with white shaft streaks;	similar to adult but most feathers have pale yellow shaft streaks;	pale yellow with grey bases to feathers;
Back:	wholly black-brown;	black-brown with some old Juvenile feathers;	brown;
Upper breast:	lanceolate black-brown with broad white shaft streaks;	similar to adults but with yellow shaft streaks;	brown with pale yellow suffusion;
Wings:	(see illustrations above);		

Juveniles undergo a partial moult soon after fledging to immature plumage that basically resembles adult plumage, but immatures retain some or all juvenile remiges and rectrices and other differences as noted above

Adult plumage is acquired early in the second year, so all adult-plumaged birds are aged (2+);

Sexing :

No difference in plumage between the sexes, but adult males are considerably larger than adult females;

Not known whether incubation is shared.