Torresian Crow Corvus orru Species No.: 692 Band size: 10 SS



Morphometrics:

The only subspecies in Australia is *C.u. cecilae* with measurements as follows:

	Adult Male	Adult Female
Wing:	315 – 382 mm	320 – 363 mm
Tail:	179 – 214 mm	176 – 221 mm
Bill:	57.5 – 66.1 mm	53.5 – 63.6 mm
Bill Depth:	19.4 – 23.8 mm	18.5 – 22.2 mm
Hackle:	23.6 – 32.7 mm	21.9 – 29.9 mm
Weight:	430 – 670 g	430 – 650 g

Ageing:

ngenig.			
Bill:	Adult (3+) black;	2 nd Immatures (2) grey- black;	1 st Immature (1) & Juvenile grey-black: initially with pinkish base
Gape:	black:	blackish-pink;	to lower mandible; pink or reddish;
•	,	• • • • • • • • • • • • • • • • • • • •	•
Palate:	black;	blackish-pink;	pink or reddish;
Interramal skin:	black;	blackish-pink;	pink;
Iris:	white, with pale blue inner ring;	hazel;	blue-grey changing to brown by three months of age;
Remiges & rectrices:	blackish with glossy greenish sheen grading	as for adult;	blackish-brown;

Juveniles undergo a partial moult to first immature plumage soon after fledging;

to bluish-purple at tips;


First Immatures retain all or most juvenile remiges, greater primary coverts, alula and rectrices. Note that outer primaries and tail feathers differ in shape from adults – see illustration below;

First immatures commence moult to second immature plumage, which is indistinguishable from adult plumage, at approximately one year old;

Second immatures, identified by subtle differences in bare parts, attain full adult colouring in a complete moult commencing at the end of their second year. Thus adults aged (3+);

Sexing:

There is no plumage dimorphism, though adult males average larger than adult females; Incubation is by the female only.



Similar species:

- The other four Australian corvid species see separate Field Data Sheets;
- · see illustrations above of the throat hackles of Australian corvids;
- see Bird in the Hand (Second Edition) Guide to Identification of Australian Corvids