
Yellow-tufted Honeyeater Lichenostomous melanops Species No.: 619 Band size: 03



Morphometrics:

	L.m. melanops (e.Great Divide)		L.m. meltoni (w. Great Divide & w. Vic)	
	Adult Male	Adult Female	Adult Male	Adult Female
THL:			38.8 – 40.4 mm	36.9 – 38.6 mm
Wing:	82 – 91 mm	78 – 88 mm	83 – 91 mm	81 – 88 mm
Wingspan:	266 – 284 mm	245 -263 mm		
Tail:	80 – 99 mm	77 -93 m	84 – 92 mm	81 – 90 mm
Weight:	23.5 – 27.0 g	21.0 – 25.5 g	23 – 26 g	21 – 23 g

 $\begin{array}{ccccc} \textit{L.m. cassidix} & (\text{Helmeted} - \text{w Gippsland}, \text{Vic}) \\ \text{THL:} & 40.0 - 43.6 \text{ mm} & 39.0 - 42.1 \text{ mm} \\ \text{Wing:} & 90 - 110 \text{ mm} & 81 - 96 \text{ mm} \\ \text{Tail:} & 90 - 110 \text{ mm} & 81 - 96 \text{ mm} \\ \text{Weight:} & 27.6 - 40.2 \text{ g} & 26.0 - 36.7 \text{ g} \end{array}$

Ageing:

tgomg.			
	Adult (2+)	Immature (2-)	Juvenile (J)
Bill:	black	horn gradually changing to black by end of second year;	greyish;
Gape:	black	yellow;	yellow;
Iris:	chestnut to brick red;	olive changing to brown;	dark brown;
Plumage:	yellow-olive	generally duller than adult;	light-brown olive;
Crown:	yellow-green	pale yellow-grey;	olive;
Back:	olive-brown tinged with brown-yellow;	dull olive-green;	light brown;
Tail feathers tip: Alula tip:	broad and pointed; pointed;	retained feathers as for juvenile; rounded;	narrow and rounded; rounded;

Juveniles moult to immature plumage within 2 months of fledging;

Retained juvenile remiges and rectrices in immatures may have synchronous growth or fault bars; Immatures moult to full adult plumage early in their second year. Thus adults are (2+).

Sexing:

Sexes similar in plumage, but males are larger than females; Female alone broods.

References: HANZAB 5; Drawings: F. Knight in HANZAB 5 - © BirdLife Australia Compiled by J.W. Hardy for the Australian Bird Study Association Inc. and reproduced with permission of BirdLife Australia.