
Red-backed Button-quail Turnix maculosa Species No.: 13 Band size: 04



Morphometrics:

 $\begin{array}{ccccc} & & \text{Adult Male} & \text{Adult Female} \\ \text{Wing:} & 72-82 \text{ mm} & 79-85 \text{ mm} \\ \text{Tail:} & 22-30 \text{ mm} & 28-33 \text{ mm} \\ \text{Weight:} & 27.5-39 \text{ g} & 38-49 \text{ g} \end{array}$

Ageing and sexing:

Post-juvenile moult within the first few months after hatching results in immature plumage. However, the illustration (reproduced above) and text in HANZAB are not consistent. The text states that the plumage of immature birds closely resemble the adults of their sex, except that they are duller in colour. Some birds can be aged on the basis of remnant juvenile feathers on breast, tertials, secondaries and primary coverts.

Moult to adult plumage commences at about 1 year of age. Thus adults are aged (2+).

Adult Male (2+) Adult Female (2+)

Bill: pale yellow with broad grey to yellow, sometimes grading to whitish-

dark grey culmen & some have yellow in distal half;

dark grey bill tip;

Supercilium: off-white faintly mottled with black- light rufous-brown; brown over ear coverts;

Lores: off-white: light rufous-brown:

Ear coverts: light brown; light brown to light rufous-brown;

Chin & throat: white; light brown;

Hindneck & upper brownish-grey; usually uniform chestnut forming a

mantle: broad chestnut collar;

Underparts: upper breast light rufous-brown rufous-brown on upper breast grading to

grading to white belly; light rufous-brown on belly.

Males do all incubation.