
Musk Lorikeet Glossopsitta concinna Species No.: 258 Band size: 24 SS



Morphometrics:

Two subspecies recognised as follows:

Nominate G.c. concinna Ssp. G.c. didimus (s-e Qld, NSW, Vic & SA) (Tasmania)

	Adult Male	Adult Female	Adult Male	Adult Female
Wing:	120 – 136 mm	117 – 134 mm	122 -136 mm	121 – 133 mm
Tail:	78 – 100 mm	83 – 96 mm	80 – 96 mm	82 – 93 mm
Bill:	11.9 – 14.4 mm	11.2 – 14.4 mm	11.8 – 12.9 mm	11.6 – 13.3 mm
Tarsus:	14.1 – 16.0 mm	14.0 – 15.7 mm	14.4 – 16.6 mm	14.0 – 16.0 mm
Weight:	60 – 90 g	57 – 86 g	58 – 72 g	61 – 73 g

Ageing:

Adult (1+) Juvenile

Bill: red to orange-yellow with narrow upper surface of upper mandible, dark

> grey-black tip, cutting edge & base; brown, rest of bill black-brown;

greyish-white & dull pink around nostrils; Cere: pale grey:

Iris: yellow-brown, orange, yellow, orangebrown;

yellow or light red; Legs & feet: brownish-grey;

light grey or straw-yellow; see below: green, some with bluish wash; Crown: Stripe behind see below; orange (feathers red with partly eve:

concealed yellow bases);

Rectrices: rounded tips; pointed tips;

Partial post-juvenile moult to adult-like immature commences at 4-5 months. Immatures retain juvenile remiges and rectrices. Moult to complete adult plumage occurs at approximately one year. Age Juveniles as (J), Immatures (1) and age Adults (1+).

Sexing:

Adult males and females are similar in size and plumage. However, females are slightly duller than males and adult birds can sometimes be sexed on subtle differences in plumage as follows:

> Adult Female Adult Male

Crown: blue: blue-green; Forehead, lores & bright red; pale red;

stripe behind eye:

Cheeks & breast; green with blue tinge; green with no blue tinge;

Female alone incubates.