


Eastern Yellow Robin *Eopsaltria australis*

Species No.: 392

Band size: 02 AY



Morphometrics:

	Nominate <i>E. a australis</i> (S.A., Vic, NSW except n.e NSW)		ssp. <i>E. a chrysorrhoea</i> (n.e. NSW & Qld)	
	Adult Male (2+)	Adult Female (2+)	Adult Male (2+)	Adult Female (2+)
THL:	37.7 – 39.3 mm	35.3 – 36.5 mm	37.6 – 41.4 mm*	36.1 – 37.3 mm*
Wing:	78 – 89 mm	76 – 87 mm	82 – 97 mm	76 – 89 mm
Wing Span:	260 – 280 mm	243 – 257 mm	265 – 285 mm *	253 -265 mm*
Tail:	60 – 70 mm	56 -67 mm	62 – 76 mm	57 – 68 mm
Weight:	16 – 30 g	14.8 – 24 g	18.5 – 27 g	15 -21.5 g

There is an intergrade zone from the Hunter River valley to the Clarence River valley.

Ageing:

- All traces of mottled brown Juvenile plumage replaced with adult-like body plumage within three months of fledging;
- Juvenile grey-brown remiges and rectrices and usually some dark brown coverts with white tips are retained by Immature.
- Immatures can be identified for up to one year by yellow soles to feet.
- Tail feathers of Immatures are pointed.
- Moults to adult plumage occurs early in the second year.
- Thus all adults 2+ and Immatures (1) in Autumn / Winter or (2-) in Spring.

Sexing :

Sexes dimorphic on THL and wing span measurements;
Female alone incubates.