
## Straw-necked lbis Threskiornis spinicollis Species No.: 180 Band size: 12 SS



## Morphometrics:

Wing: 364 – 415 mm 351 – 375 r	
8 <sup>th</sup> primary: 227 – 246 mm 215 – 237 r	nm
Tail: 130 – 152 mm 126 – 142 r	nm
Bill: 156 – 185 mm 120 – 142.2	2 mm
Tarsus: 83.6 – 102 mm 75.4 – 83.2	mm
Toe: 75.2 – 87.2 mm 69.7 – 93.6	mm
Weight: 1400 – 1570 g 1150 – 132	0 g

## Ageing & Sexing:

Timing and duration of moults is not known, so age of an immatures first attaining adult plumage and first breeding is not known. However, there are juvenile and immature plumage phases and adult plumage is attained gradually by balding and colouring of plumage. Thus adults will be at least in their second year.

Differences between plumage and bare parts of adults and juveniles /Immatures are:

Dinoronoco b	Adult Male	Adult Female	Juvenile/Immature
Bill:	blackish grey with pale olive to blackish rugose plates;		fuscous black to black;
Palate:	back – dark grey, front – olive-grey;	dark grey;	
Iris:	dark brown;	dark brown;	umber - turning brown;
Head & foreneck:	bare grey-black skin;	greyish-black skin;	black-brown to dark brown;
Hindneck:	dark brown with glossy dark green tips;	as for male;	dark brown;
Upper breast	glossy dark green, basally dark brown forming incomplete band across breast centrally white;	complete wide dark band of feathers;	upper breast dark brown
Upper tibia:	light pink with black lines;	bright red-pink;	blackish grey;
Rest of legs:	blackish red;	dull greyish-red;	blackish grey;

Adult males have a definitively longer bill than adult females; Both sexes incubate.