

### ACKNOWLEDGMENTS

Data used in this paper were collected for a M. Nat. Res. thesis in the University of New England. I thank my supervisors J. S. Weir and P. J. Jarman for their encouragement and guidance. My thanks to R. Sharrad and T. C. R. White for comments on the manuscript and to R. Roberts for preparing the figures.

### REFERENCES

- Beckle, H. (1983). Effects of unseasonable rains in January 1982 on waterfowl in south-western Australia. 1. Responses of selected species on coastal summer refuges. *West Aust. Nat.* 15: 117-122.
- Briggs, S. V. (1977). Variations in waterbird numbers at four swamps on the Northern Tablelands of New South Wales. *Aust. Wildl. Res.* 4: 301-309.
- Briggs, S. V. (1979). Daytime habitats of waterbirds at four swamps on the Northern Tablelands of New South Wales. *Emu* 79: 211-214.
- Carrick, R. (1962). Breeding, movement, and conservation of ibises (Threskiornidae) in Australia. *CSIRO Wildl. Res.* 7: 71-88.
- Ford, J. R. (1958). Seasonal variations in populations of anatidae at the Bibra Lake District, Western Australia. *Emu* 58: 31-41.
- Frith, H. J. (1982). 'Waterfowl in Australia' (Angus and Robertson: Sydney.)
- Gentilli, J. and Beckle, H. (1983). Modelling a climatically pulsating population: Grey Teal in South-western Australia. *J. Biogeography* 10: 75-96.
- Gosper, D. G., Briggs, S. V. and Carpenter, S. M. (1983). Waterbird dynamics in the Richmond Valley, New South Wales, 1974-77. *Aust. Wildl. Res.* 10: 319-327.
- Harper, M. J. (1990). Waterbird dynamics at Bool Lagoon, South Australia, 1983-87. *Aust. Wildl. Res.* 17: 113-122.
- Hnatiuk, S. H. (1987). Numbers of waterbirds at a swamp in Ferndale, Western Australia, 1978 to 1981. *Corella* 11: 20-27.
- Lavery, H. J. (1970). Studies of waterfowl in north Queensland. 4. Movements. *Aust. J. Agric. and Anim. Sci.* 27: 411-424.
- Norman, F. I. (1983). Grey Teal, Chestnut Teal, and Pacific Black Duck at a saline habitat in Victoria. *Emu* 83: 262-271.
- Norman F. I. and Nicholls, L. (1991). The Southern Oscillation and variations in waterfowl abundance in southeastern Australia. *Aust. J. Ecol.* 16: 485-490.
- Ryan, T. A. Jr., Joiner, B. L. and Ryan, B. F. (1976). 'Minitab Student Handbook.' (Duxburg Press: Boston.)
- Sharland, M. (1960). The Tasmanian coot mystery. *Emu* 60: 95-98.
- White, J. M. (1986a). Breeding of Black Swans on two New England lagoons. *Corella* 10: 17-20.
- White, J. M. (1986b). The management of the New England Lagoons for waterbirds. M. Nat. Res. thesis. University of New England.
- Whyte, R. J. (1981). Winter fluctuations in waterbird numbers on a northern tablelands lagoon of New South Wales. *Emu* 81: 243-246.
- Woodall, P. F. (1985). Waterbird populations in the Brisbane Region, 1972-83 and correlates with rainfall and water heights. *Aust. Wildl. Res.* 12: 495-506.

---

### ERRATUM

#### Corella 14(3)

Brown, R. J., Brown, M. N. and Russell, E. M. (1990). The survival of four species of passerine in Karri forests in south-western Australia. *Corella* 14(3): 69-78.

**1. In Appendix 1, page 78, line 6.**

Years 1-7 should read Years 1-8.

**2. on pages 75 and 76.**

The mean annual survival rate of the Red-winged Fairy Wren for years 1-8 after banding, derived from Table 6, should be 78% and not 67%.