Black-tailed Native-hen Gallinula ventralis

Species Code: 344 Band: [T elliptical]



Morphometrics:

	Adult Male	Adult Female
Wing:	205 – 231 mm	200 – 233 mm
8 th Primary:	139 – 159 mm	131 – 160 mm
Tail:	76 – 88 mm	72 – 87 mm
Tarsus:	59 – 66 mm	50.5 – 62.4 mm
Weight:	250 – 530 g	322 – 405 g

Ageing:

	Adult (2+) & immature (1)	Juvenile (J)
Bill & frontal	light green, dull green or pale bluish-	greenish yellow with dusky tip;
ماما مام	anno esta de la comunidad de la Danad half	

shield: grey tip to lower mandible. Basal half of lower mandible orange red to red;

Iris: orange-yellow or sometimes yellow;

Legs & feet: tarsus & toes pinkish red and claws

grev-black:

adults sexually dimorphic (see below); Plumage:

۱۱ مانوم، ۱۱

black-brown;

dirty pink with brown centres to scutes;

similar to adult female, but duller and feathers generally smaller & narrower; flank spots much smaller than adults.

Juvenile moult to immature plumage within a few months of leaving nest, but immatures retain some juvenile feathers. Adult plumage attained in moult coinciding with adult post-breeding moult; commencing at approximately 1 year old. Thus adults are aged (2+).

Sexing:

Flanks:

Adult males tend to be larger than females (see above); Adults are sexually dimorphic on plumage (see below):

Adult Male

Base of bill: black-brown feathers form dark band; Crown to hindneck: olive-brown to dark olive-brown;

Upperparts: uniform olive;

Breast & belly: dark grey or dark leaden grey; lighter and greyer than male; pale olive to olive-grey; leaden grey to grey often with

Adult Female

band narrower or absent:

bluish tinge;

white spots smaller than male.

No information on incubation roles.

large elongated white tips to feathers;

ABSA – Bird in the Hand (Second Edition) – New Zealand species	s. Published on www.absa.asn.au – January 2022